



## Home Work 1

1. Figure *a* shows an arrangement of three charged particles separated by distance  $d$ . Particles *A* and *C* are fixed on the  $x$  axis, but particle *B* can be moved along a circle centered on particle *A*. During the movement, a radial line between *A* and *B* makes an angle  $\theta$  relative to the positive direction of the  $x$  axis (Fig. *b*). The curves in Fig. *c* give, for two situations, the magnitude  $F_{\text{net}}$  of the net electrostatic force on particle *A* due to the other particles. That net force is given as a function of angle  $\theta$  and as a multiple of a basic amount  $F_0$ . For example on curve 1, at  $\theta = 180^\circ$ , we see that  $F_{\text{net}} = 2F_0$ . (a) For the situation corresponding to curve 1, what is the ratio of the charge of particle *C* to that of particle *B* (including sign)? (b) For the situation corresponding to curve 2, what is that ratio?



2. In crystals of the salt cesium chloride, cesium ions  $\text{Cs}^+$  form the eight corners of a cube and a chlorine ion  $\text{Cl}^-$  is at the cube's center (see the following figure). The edge length of the cube is 0.40 nm. The  $\text{Cs}^+$  ions are each deficient by one electron (and thus each has a charge of  $+e$ ), and the  $\text{Cl}^-$  ion has one excess electron (and thus has a charge of  $-e$ ). (a) What is the magnitude of the net electrostatic force exerted on the  $\text{Cl}^-$  ion by the eight  $\text{Cs}^+$  ions at the corners of the cube? (b) If one of the  $\text{Cs}^+$  ions is missing, the crystal is said to have a *defect*; what is the magnitude of the net electrostatic force exerted on the  $\text{Cl}^-$  ion by the seven remaining  $\text{Cs}^+$  ions?



3. What equal positive charges would have to be placed on Earth and on the Moon to neutralize their gravitational attraction? (b) Why don't you need to know the lunar distance to solve this problem? (c) How many kilograms of hydrogen ions (that is, protons) would be needed to provide the positive charge calculated in (a)?
4. A nonconducting spherical shell, with an inner radius of 4.0 cm and an outer radius of 6.0 cm, has charge spread nonuniformly through its volume between its inner and outer surfaces. The *volume charge density*  $\rho$  is the charge per unit volume, with the unit coulomb per cubic meter. For this shell  $\rho = b/r$ , where  $r$  is the distance in meters from the center of the shell and  $b = 3.0 \mu\text{C}/\text{m}^2$ . What is the net charge in the shell?
5. Earth's atmosphere is constantly bombarded by *cosmic ray protons* that originate somewhere in space. If the protons all passed through the atmosphere, each square meter of Earth's surface would intercept protons at the average rate of 1500 protons per second. What would be the electric current intercepted by the total surface area of the planet?