Home Work 12

12-1 In Fig.1, an electric field is directed out of the page within a circular region of radius R = 3.00 cm. The field magnitude is $E = (0.500 \text{ V/m} \cdot \text{s})(1 - r/R)t$, where *t* is in seconds and *r* is the radial distance ($r \le R$). What is the magnitude of the induced magnetic field at radial distances (a) 2.00 cm and (b) 5.00 cm?

12-2 In Fig. 2, a parallel-plate capacitor has square plates of edge length L = 1.0 m. A current of 2.0 A charges the capacitor, producing a uniform electric field \overrightarrow{E} between the plates, with \overrightarrow{E} perpendicular to the plates. (a) What is the displacement current i_d through the region between the plates? (b) What is dE/dt in this region? (c) What is the displacement current encircled by the square dashed path of edge length d = 0.50 m? (d) What is $\oint \overrightarrow{B} \cdot d\overrightarrow{s}$ around this square dashed path?

12-3 Assume that an electron of mass m and charge magnitude e moves in a circular orbit of radius r about a nucleus. A uniform magnetic field \overrightarrow{B} is then established perpendicular to the plane of the orbit. Assuming also that the radius of the orbit does not change and that the change in the speed of the electron due to field \overrightarrow{B} is small, show that the change in the orbital magnetic dipole moment of

the electron due to the field is
$$\Delta \mu = \frac{e^2 r^2 B}{4m_e}$$
.

12-4 The magnetic field of Earth can be approximated as the magnetic field of a dipole. The horizontal and vertical components of this field at any distance r from Earth's center are given by

$$B_h = \frac{\mu_0 \mu}{4\pi r^3} \cos \lambda_m, \qquad B_v = \frac{\mu_0 \mu}{2\pi r^3} \sin \lambda_m,$$

where λ_m is the magnetic latitude (this type of latitude is measured from the geomagnetic equator toward the north or south geomagnetic pole). Assume that Earth's magnetic dipole moment has magnitude $\mu = 8.00 \times 10^{22} \text{ A} \cdot \text{m}^2$. (a) Show that the magnitude of Earth's field at latitude λ_m is

given by $B = \frac{\mu_0 \mu}{4\pi r^3} \sqrt{1 + 3\sin^2 \lambda_m}.$

(b) Show that the inclination φ_i of the magnetic field is related to the magnetic latitude λ_m by . $\tan \phi_i = 2 \tan \lambda_m$.

12-5 A charge q is distributed uniformly around a thin ring of radius r. The ring is rotating about an axis through its center and perpendicular to its plane, at an angular speed ω . (a) Show that the magnetic moment due to the rotating charge has magnitude (b) What is the direction of this magnetic $\mu = \frac{1}{2}q\omega r^2$ moment if the charge is positive?

12-6 Consider a solid containing N atoms per unit volume, each atom having a magnetic dipole moment μ . Suppose the direction of μ can be only parallel or antiparallel to an externally applied magnetic field *B* (this will be the case if μ is due to the spin of a single electron). According to statistical mechanics, the probability of an atom being in a state with energy *U* is proportional to $e^{-U/kT}$, where *T* is the temperature and *k* is Boltzmann's constant. Thus, because energy *U* is - $\mu \cdot B$, the fraction of atoms whose dipole moment is parallel to *B* is proportional to $e^{-\mu B/kT}$ and the fraction of atoms whose dipole moment is antiparallel to is proportional to $e^{-\mu B/kT}$. (a) Show that the magnitude of the magnetization of this solid is $M = N \mu \tanh(\mu B/kT)$. Here tanh is the hyperbolic tangent function: $\tanh(x) = (e^x - e^{-x})/(e^x + e^{-x})$. (b) Show that the result given in (a) reduces to $M = N \mu^2 B/kT$ for $\mu B < < kT$. (c) Show that the result of (a) reduces to $M = N \mu$ for $\mu B >> kT$. (d) Show that both (b) and (c) agree qualitatively with Fig. 3.

